

06-28-00

A

Please type a plus sign (+) inside this box → ☐

PTO/SB/05 (4/98)
Approved for use through 09/30/2000. OMB 0651-0032
Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

06/27/00
JC712 U.S. PTO

UTILITY PATENT APPLICATION TRANSMITTAL

(Only for new nonprovisional applications under 37 C.F.R. § 1.53(b))

Attorney Docket No.	GI-5238BA
First Inventor or Application Identifier	Nicholas W. Warne
Title	Topical Formulation for Delivery of Interleukin-11
Express Mail Label No.	EL564796563US

APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO: Assistant Commissioner for Patents
Box Patent Application
Washington, DC 20231

- ☒ * Fee Transmittal Form (e.g., PTO/SB/17)
(Submit an original and a duplicate for fee processing)
- ☒ Specification [Total Pages **34**]
(preferred arrangement set forth below)
 - Descriptive title of the Invention
 - Cross References to Related Applications
 - Statement Regarding Fed sponsored R & D
 - Reference to Microfiche Appendix
 - Background of the Invention
 - Brief Summary of the Invention
 - Brief Description of the Drawings (if filed)
 - Detailed Description
 - Claim(s)
 - Abstract of the Disclosure
- ☐ Drawing(s) (35 U.S.C. 113) [Total Sheets ☐
- Oath or Declaration [Total Pages **3**]
 - ☐ Newly executed (original or copy)
 - ☒ Copy from a prior application (37 C.F.R. § 1.63(d))
(for continuation/divisional with Box 16 completed)
 - ☐ DELETION OF INVENTOR(S)
Signed statement attached deleting inventor(s) named in the prior application, see 37 C.F.R. §§ 1.63(d)(2) and 1.33(b).

- ☐ Microfiche Computer Program (Appendix)
- Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)
 - ☐ Computer Readable Copy
 - ☐ Paper Copy (identical to computer copy)
 - ☐ Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

- ☐ Assignment Papers (cover sheet & document(s))
- ☐ 37 C.F.R. §3.73(b) Statement ☐ Power of Attorney
(when there is an assignee)
- ☐ English Translation Document (if applicable)
- ☐ Information Disclosure Statement (IDS)/PTO-1449 ☐ Copies of IDS Citations
- ☐ Preliminary Amendment
- ☐ Return Receipt Postcard (MPEP 503)
(Should be specifically itemized)
- ☐ * Small Entity Statement filed in prior application, Status still proper and desired
(PTO/SB/09-12)
- ☐ Certified Copy of Priority Document(s)
(if foreign priority is claimed)
- ☐ Other: _____

* NOTE FOR ITEMS 1 & 13: IN ORDER TO BE ENTITLED TO PAY SMALL ENTITY FEES, A SMALL ENTITY STATEMENT IS REQUIRED (37 C.F.R. § 1.27), EXCEPT IF ONE FILED IN A PRIOR APPLICATION IS RELIED UPON (37 C.F.R. § 1.28).


16. If a CONTINUING APPLICATION, check appropriate box, and supply the requisite information below and in a preliminary amendment:

<input checked="" type="checkbox"/> Continuation	<input type="checkbox"/> Divisional	<input type="checkbox"/> Continuation-in-part (CIP)	of prior application No: 09 / 179,026
Prior application information: Examiner P. Mertz			Group / Art Unit: 1646

For CONTINUATION or DIVISIONAL APPS only: The entire disclosure of the prior application, from which an oath or declaration is supplied under Box 4b, is considered a part of the disclosure of the accompanying continuation or divisional application and is hereby incorporated by reference. The incorporation can only be relied upon when a portion has been inadvertently omitted from the submitted application parts.

17. <input checked="" type="checkbox"/> Customer Number or Bar Code Label (Insert C)	 24363	18. <input type="checkbox"/> Correspondence address below (e label here)
---	---	---

Name	M. Kymne Hehman				
	American Home Products Corporation				
Address	Patent & Trademark Office -2B (Attn: Kay E. Brady)				
	One Campus Drive				
City	Parsippany	State	New Jersey	Zip Code	07054
Country		Telephone		Fax	

Name (Print/Type)	M. Kymne Hehman	Registration No. (Attorney/Agent)	39,206
Signature		Date	27 June 2000

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231.

-1-

Date: 6/27/00 Express Mail Label No. EL564 7965634S

Inventor: Nicholas W. Warne, Camille L. Bedrosian, James C.
Keith, Jr., Ullrich S. Schwertschlag and Paul F. Schendal
Attorney's Docket No.: GI-5238BA

TOPICAL FORMULATION FOR DELIVERY OF INTERLEUKIN-11

RELATED APPLICATION(S)

This is a Continuation of U.S. Application No. 09/179,026, filed October 26, 1998, which is a continuation-in-part of U.S. Application No. 08/892,407, filed July 15, 1997, which is a divisional of U.S. Application No. 08/495,724, filed June 27, 1995, now Patent No. 5,679,339, issued October 21, 1997, the entire teachings of which are incorporated herein by reference.

FIELD OF INVENTION

The present invention relates generally to novel compositions and methods for topical delivery of interleukin-11 (IL-11). In preferred embodiments, patients are treated employing topical delivery of recombinant human IL-11 for inflammatory bowel diseases (e.g., Crohn's disease, ulcerative colitis, indeterminate colitis, and infectious colitis), mucositis (e.g., oral mucositis, gastrointestinal mucositis, nasal mucositis, and proctitis), necrotizing enterocolitis, inflammatory skin disorders (e.g., psoriasis, atopic dermatitis, and contact hypersensitivity), aphthous ulcers, pharyngitis, esophagitis, peptic ulcers, gingivitis, periodontitis, and ocular diseases (e.g., conjunctivitis, retinitis, and uveitis).

BACKGROUND OF THE INVENTION

Inflammatory responses include a broad range of host reaction to a variety of insults, such as injury, infection, or rejection. It is the overproduction of mediators that is believed to be associated with a broad range of disorders, including AIDS, arthritis (rheumatoid arthritis, osteoarthritis, spondyloarthropathies), antibiotic-induced diarrheal diseases, multiple sclerosis, osteoporosis, gingivitis, peptic ulcer disease, esophagitis, diabetes, retinitis, uveitis, reperfusion injury after myocardial infarction, cerebral vascular accident, aphthous ulcers (oral), atherosclerosis, tumor metastases, asthma, preeclampsia, pancreatitis, psoriasis, infertility and allergic disorders such as rhinitis, conjunctivitis, and urticaria.

Some of these disorders and their symptoms are briefly summarized below. According to the methods of the present invention, IL-11 is administered topically to modulate the host's over reaction at the site of insult, thereby treating the following disorders.

Inflammatory Bowel Disease: Inflammatory bowel disease is the term generally applied to four diseases of the bowel, namely Crohn's disease, ulcerative colitis, indeterminate colitis, and infectious colitis.

Ulcerative Colitis: Ulcerative colitis is a chronic inflammatory disease of unknown etiology afflicting the large intestine. The course of the disease may be continuous or relapsing, mild or severe. The earliest lesion is an inflammatory infiltration with abscess formation at the base of the crypts of Lieberkühn. Coalescence of these distended and ruptured crypts tends to separate the overlying mucosa from its blood supply, leading to ulceration. Signs and symptoms of the disease include cramping, lower abdominal pain, rectal bleeding, and frequent, loose discharges consisting mainly of blood, pus, and mucus with scanty fecal particles. A total colectomy may be required for acute severe or chronic, unremitting ulcerative colitis.

Crohn's Disease: Crohn's disease (also known as regional enteritis or ulcerative ileitis) is also a chronic inflammatory disease of unknown etiology but, unlike

ulcerative colitis, it can affect any part of the bowel. The most prominent feature of the disease is the granular, reddish-purple edematous thickening of the bowel wall. With the development of inflammation, these granulomas often lose their circumscribed borders and integrate with the surrounding tissue. Diarrhea and obstruction of the bowel are the predominant clinical features. As with ulcerative colitis, the course of the disease may be continuous or relapsing, mild or severe but, unlike ulcerative colitis, it is not curable by resection of the involved segment of bowel. Most patients with Crohn's disease require surgery at some point, but subsequent relapse is common and continuous medical treatment is usual.

- 10 **Mucositis:** Mucositis involves ulcerative breakdown of mucosal epithelial tissue, and is literally defined as inflammation of the mucous membrane. The pathophysiology of mucositis in response to toxic insults to the mucosa by chemotherapy or by ionizing radiation is complex and involves a cascade of interactions among cells, cytokines and the oral microflora. The underlying premise for
- 15 susceptibility of the mucosa of the oropharynx and gastrointestinal tract to chemotherapy or radiation damage is related to rapid epithelial stem cell turnover. Mucositis may be characterized by the following phases:

1. Early inflammatory phase characterized by release of inflammatory cytokines in response to local tissue damage caused by cytotoxic agent(s);
- 20 2. Epithelial phase characterized by death of basal cells, which hinders re-population of the epithelium. This inability to regenerate leads to atrophy followed by ulceration. The ulceration represents loss of an important anatomic barrier at a site of local microflora;
3. Infection phase characterized by local invasion of microflora that results
- 25 in an inflammatory response to the local infection. The inflammation results in additional local tissue damage and possibly erosive ulceration; and
4. Healing phase characterized by resolution of the infection and regeneration of epithelium.

Oral mucositis produces the following clinical symptoms and signs resulting from cellular damage: 1) sensation of dryness; 2) asymptomatic redness and erythema; 3) solitary white elevated desquamative patches which are painful upon pressure contact; and 4) large, painful, contiguous pseudomembranous lesions associated with dysphagia and decreased oral intake. These spontaneously painful lesions histopathologically show loss of epithelial cells to the basement membrane, which exposes the connective tissue stroma with its associated innervation.

As with oral mucosa, gastrointestinal mucosal damage results from disturbance of cellular mitosis that leads to reduction in the turnover rate of the basal cells of the intestinal crypts. The symptoms and signs of gastrointestinal mucositis include tenesmus (painful ineffectual straining at stool), pain, bleeding, diarrhea, telangiectasia (neovascularization), and progression to ulceration. Early signs of diarrhea include increased stool frequency, loose or watery stool, food aversion, increased bowel sounds, abdominal pain, and some loss of skin turgor indicative of dehydration. When the diarrhea is severe it may be associated with mucosal ulceration, bleeding, intestinal perforation and proctitis. Stool exam may reveal occult blood and fecal leukocytes.

Necrotizing Enterocolitis: Necrotizing enterocolitis is an inflammatory disease of unknown etiology that afflicts between 1-5% of all infants admitted to neonatal intensive care units, most of whom are premature infants. Signs and symptoms include abdominal distention, gastrointestinal hemorrhage, and feeding intolerance. The disease most often involves the ileum and colon, and is characterized by loss of epithelium and submucosal edema, ulcerations, and, in severe cases, transmural necrosis.

Psoriasis: Psoriasis is a chronic inflammatory skin disorder involving hyperproliferation of the epidermis and inflammation of both the epidermis and the dermis. Macrophage, T-cells and neutrophil infiltration of the dermis and epidermis is seen, and proinflammatory mediators are released from these activated cells.

Aphthous Ulcers (oral): Although the cause of aphthous ulcers remain unknown, many physicians believe they are caused by autoimmune phenomena, which cause the destruction of discrete areas of the oral mucosa which leads to oral ulceration. Among the cytokines present in these active areas of ulceration, TNF- α appears to play
5 a predominant role.

Gingivitis: Adult periodontitis is strongly associated with infection by *Porphyromonas gingivalis*. Proteolytic enzymes, which are produced in large quantity by this bacteria, are considered as important pathogenic agents. The increased production and flow of gingival crevicular fluid (GCF) is an important change in
10 gingival tissues during periodontal infection, correlating with clinical indices of gingival inflammation. Indeed, salivary protein and albumin concentrations of individuals with periodontitis, which are an indication of plasma leakage due to vascular permeability enhancement (VPE), are significantly increased compared to healthy subjects. The production of GCF appears dependent on VPE induced at periodontitis sites,
15 presumably involving proteinase(s) of *P. gingivalis* in their generation.

Esophagitis: The most common cause of esophagitis is the chronic reflux of hydrochloric acid from the stomach due to inefficiency of the cardiac sphincter of the stomach. The chronic presence of acid in the lower esophagus leads to damage of the esophageal mucosa. In the most severe form, a syndrome called Barrett's esophagus
20 can develop which often leads to esophageal cancer. Other causes of esophagitis include parenteral chemotherapy and ionizing radiation, associated with radiation therapy for cancer in the thoracic cavity.

Retinitis: Inflammation of the light sensitive retina, retinitis, can occur due to a variety of viral, bacterial or autoimmune etiologies. The end result is destruction of the
25 retina and loss of sight.

Uveitis: Inflammation of the anterior portion of the eye its associated structures, the iris and cornea occurs with a relatively high frequency in patients with autoimmune disorders.

Peptic Ulcer Disease: Inhibition of gastric acid secretion with H₂-receptor antagonists and, more recently, blockers of H⁺,K⁺-ATPase (also known as the proton pump) has been the mainstay of therapy for peptic ulcer disease. The pathophysiology of peptic ulcers remains obscure. An appreciation of the complexity of the physiology of the gastric mucosa has led to a hypothesis that peptic ulcers are the result of an imbalance in the relative importance of aggressive (acid, pepsin) and protective (mucus, bicarbonate, blood flow, prostaglandins, *etc.*) factors. Infection of the mucosa of the human gastric antrum with the bacterium *Helicobacter pylori* has been widely accepted as the cause of chronic, active, type B gastritis. Further, this form of gastritis has been linked directly to peptic ulcer disease by studies showing that eradication of *H. pylori* reverses this gastritis and prevents duodenal ulcer relapse. Because cytokines are the principal mediators by which immune/inflammatory cells communicate with each other and with other cells, it is likely that these small peptides are involved in the pathogenesis of chronic active type B gastritis and the resulting peptic ulcer disease.

Some cytokines (IL-1, epidermal growth factor, transforming growth factor- α , acidic and basic fibroblast growth factors) tip the balance towards peptic ulcer healing; others (tumor necrosis factor- α) appear to have no effect; still others (IL-4) may even cause gastrointestinal damage.

The infiltration and activation of multiple types of inflammatory cells result in a series of degenerative changes in the vasculature of the affected area, as well as inciting damage of the surrounding parenchymal tissue.

SUMMARY OF THE INVENTION

Provided by the present invention are novel compositions for the topical administration of IL-11 in order to treat a number of disorders where such administration is preferable to systemic administration. Among the reasons that local administration may be preferred are the ease of administering a topical formulation compared to administration of subcutaneous injectable formulations. In certain classes

of patients, the toxicity profile of chemotherapeutic agents may be such that concurrent parenteral administration of IL-11 is relatively unsuited. Other patients may have medical conditions for which the adverse event profile of parenteral IL-11 is relatively unsuited.

- 5 According to the present invention, IL-11, analogs, and derivatives thereof, are administered to patients, either prophylactically or at the onset of symptoms associated with the aforementioned disorders, through one of several forms of local administration described below. IL-11 can be administered in suitable pharmaceutically acceptable carriers either alone or in combination with other conventional agents useful in
- 10 alleviating the symptoms associated with the aforementioned disorders.

- In one embodiment, the present invention comprises preparations of IL-11 which are suitable for oral delivery to the mouth. Suitable oral preparations may be prepared with aqueous-based solutions such as sodium bicarbonate (e.g., Brioschi®), or in gels and suspensions for topical administration in the mouth. Oral preparations may
- 15 also take the form of patches for delivery of IL-11 to the mouth via sustained release. Additional oral preparations may comprise IL-11 in the form of a lozenge or an uncoated tablet which is retained in the mouth. The oral preparations are particularly well-suited for disorders and inflammatory responses involving the mucosa of the head, neck and/or mouth. Such conditions include oral mucositis, pharyngitis, esophagitis,
- 20 gingivitis, periodontitis, and aphthous ulcers (oral). Such conditions may result, for example, from chemotherapy or radiotherapy for head and neck cancer, cervical esophageal cancer or lung cancer.

- In other embodiments, the present invention comprises preparations of IL-11 which are suitable for topical delivery for mucosa and/or dermis. Such topical
- 25 preparations may be prepared in the form of aqueous-based solutions, gels, ointments or creams for topical administration, as gels and suspensions for cervical administration, as pills, tablets, capsules or suppositories for immediate or sustained release to the gastrointestinal tract, or in the form of solution for enema. Such topical preparations are

especially suited for treatment of disorders relating to local regions, such as psoriasis, as well as inflammatory bowel disease, esophagitis, and gastrointestinal mucositis. Some of these conditions may result, for example, from chemotherapy and/or radiotherapy for colorectal cancer, prostate cancer, cervical esophageal cancer or lung cancer.

- 5 Suitable doses of IL-11 are generally in the range of between about 1 and about 250 $\mu\text{g/kg}$ body weight, and may be administered from once a week up to about six times daily. Treatment may continue for a period of between one day and six months, or for as long as is deemed necessary and safe in the treatment of the aforementioned disorders, as is readily ascertained by standard tests by the attending physician,
- 10 depending upon the nature of the disorder being treated.

DETAILED DESCRIPTION OF THE INVENTION

- Provided by the present invention are methods and compositions for topical administration of IL-11, for the treatment of various disorders, such as inflammatory bowel diseases (e.g., Crohn's disease, ulcerative colitis, indeterminate colitis, and
- 15 infectious colitis), mucositis (e.g., oral mucositis, gastrointestinal mucositis, nasal mucositis, and proctitis), necrotizing enterocolitis, aphthous ulcers, psoriasis, pharyngitis, esophagitis, peptic ulcers, gingivitis, periodontitis, and ocular diseases (e.g., conjunctivitis, retinitis, and uveitis).

- Unless defined otherwise, all technical and scientific terms used herein have the
- 20 same meaning as commonly understood by one of ordinary skill in the art to which this invention pertains. Although any methods and materials similar or equivalent to those described herein can be used in the practice of the present invention, the preferred methods and materials are described. For purposes of the present invention, the following terms are defined below.

- 25 As used herein, the term "topical" refers to application at the site of inflammation (i.e., non-systemic administration), and includes, for example, oral, intranasal, intrabronchial, topical, and rectal routes of administration. The term "topical

formulation” refers to a pharmaceutical formulation which is suitable for topical administration of the active ingredient.

As used herein, the term “pharmaceutically acceptable” refers to a carrier medium which does not interfere with the effectiveness of the biological activity of the active ingredient(s) and which is not toxic to the host to which it is administered.

As used herein, the terms “therapeutically effective amount” and “therapeutically effective dose” as applied to the active ingredient refers to the amount of the component in the composition or administered to the host that results in an increase in the therapeutic index of the host. The “therapeutic index” can be defined for purposes herein in terms of efficacy, i.e., extent of reduction or inhibition of inflammation. Suitable doses of the active ingredient can be determined using well-known methods, a variety of which are known and readily available in the pharmaceutical sciences, including, for example, measurement of markers associated with the disorder (e.g., TNF- α , IL-1 β , IL-6, and/or IL-12 p40 mRNA), the biological effects of TNF- α , and decreased symptomatology.

Interleukin 11 (IL-11) is a pleiotropic cytokine that stimulates primitive lymphohematopoietic progenitor cells and synergizes with other hematopoietic growth factors to stimulate the proliferation and maturation of megakaryocytes. IL-11 is described in detail in International Application PCT/US90/06803, published May 30, 1991, as well as in U.S. Patent No. 5,215,895; issued June 1, 1993. A cloned human IL-11 was previously deposited with the ATCC, 12301 Parklawn Drive, Rockville, Maryland, on March 30, 1990 under ATCC No. 68284. Moreover, as described in U.S. Patent No. 5,270,181, issued December 14, 1993, and U.S. Patent No. 5,292,646, issued March 8, 1994, IL-11 may also be produced recombinantly as a fusion protein with another protein. IL-11 can be produced in a variety of host cells by resort to now conventional genetic engineering techniques. In addition, IL-11 can be obtained from various cell lines, for example, the human lung fibroblast cell line, MRC-5 (ATCC Accession No. CCL 171), and Paul *et al.*, the human trophoblastic cell line, TPA30-1

(ATCC Accession No. CRL 1583). A cDNA encoding human IL-11, as well as the deduced amino acid sequence (amino acids 1 to 199), is described in *Proc. Natl. Acad. Sci. USA* 87:7512 (1990). U.S. Patent No. 5,292,646, *supra*, describes a des-Pro form of IL-11 in which the N-terminal proline of the mature form of IL-11 (amino acids 22-199) has been removed (amino acids 23-199). As is appreciated by one skilled in the art, any form of IL-11 which retains IL-11 activity, such as variants through the substitution or deletion of amino acids, analogs and derivatives of IL-11, is useful according to the present invention. The disclosure of each of the above publications is hereby incorporated by reference for the contents thereof.

10 In addition to recombinant techniques, IL-11 may also be produced by known conventional chemical synthesis. Methods for constructing the polypeptides useful in the present invention by synthetic means are known to those of skill in the art. The synthetically constructed cytokine polypeptide sequences, by virtue of sharing primary, secondary, or tertiary structural and conformational characteristics with the natural
15 cytokine polypeptides are anticipated to possess biological activities in common therewith. Such synthetically constructed cytokine polypeptide sequences or fragments thereof, which duplicate or partially duplicate the functionality thereof may also be used in the method of this invention. Thus, they may be employed as biologically active or immunological substitutes for the natural, purified cytokines useful in the present
20 invention.

Modifications in the protein, peptide or DNA sequences of these cytokines or active fragments thereof may also produce proteins which may be employed in the methods of this invention. Such modified cytokines can be made by one skilled in the art using known techniques. Modifications of interest in the cytokine sequences, *e.g.*,
25 the IL-11 sequence, may include the replacement, insertion or deletion of one or more selected amino acid residues in the coding sequences. Mutagenic techniques for such replacement, insertion or deletion are well known to one skilled in the art. (See, *e.g.*, U. S. Patent No. 4,518,584.)

Other specific mutations of the sequences of the cytokine polypeptides which may be useful therapeutically as described herein may involve, *e.g.*, the insertion of one or more glycosylation sites. An asparagine-linked glycosylation recognition site can be inserted into the sequence by the deletion, substitution or addition of amino acids into the peptide sequence or nucleotides into the DNA sequence. Such changes may be made at any site of the molecule that is modified by addition of O-linked carbohydrate. Expression of such altered nucleotide or peptide sequences produces variants which may be glycosylated at those sites.

Additional analogs and derivatives of the sequence of the selected cytokine which would be expected to retain or prolong its activity in whole or in part, and which are expected to be useful in the present method, may also be easily made by one of skill in the art. One such modification may be the attachment of polyethylene glycol (PEG) onto existing lysine residues in the cytokine sequence or the insertion of one or more lysine residues or other amino acid residues that can react with PEG or PEG derivatives into the sequence by conventional techniques to enable the attachment of PEG moieties.

Additional analogs of these selected cytokines may also be characterized by allelic variations in the DNA sequences encoding them, or induced variations in the DNA sequences encoding them. It is anticipated that all analogs disclosed in the above-referenced publications, including those characterized by DNA sequences capable of hybridizing to the disclosed cytokine sequences under stringent hybridization conditions or non-stringent conditions (Sambrook *et al.*, Molecular Cloning. A Laboratory Manual, 2d edit., Cold Spring Harbor Laboratory, New York (1989)) will be similarly useful in this invention.

Also considered as derivatives useful in these methods are fusion molecules, prepared by fusing the sequence or a biologically active fragment of the sequence of one cytokine to another cytokine or proteinaceous therapeutic agent, *e.g.*, IL-11 fused to IL-6 (see, *e.g.*, methods for fusion described in PCT/US91/06186 (WO92/04455),

published March 19, 1992). Alternatively, combinations of the cytokines may be administered together according to the method.

Thus, where in the description of the methods of this invention IL-11 is mentioned by name, it is understood by those of skill in the art that IL-11 encompasses
5 the protein produced by the sequences presently disclosed in the art, as well as proteins characterized by the modifications described above yet which retain substantially similar activity. Standard laboratory tests are utilized to monitor progress of the treatment. Levels of TNF- α in serum or the biologic effects of TNF- α could be followed in a variety of these diseases. Decreased symptomatology could also be used
10 to monitor the effectiveness of treatment as is well known to physicians skilled in the art of treating such disorders. Treatment is preferably prophylactic, but may also be at the onset of symptoms associated with the aforementioned disorders.

In one aspect of the invention, IL-11 is used in a therapeutic composition to alleviate or prevent the onset of symptoms associated with an inflammatory disorder.
15 While it is possible to administer IL-11 alone, it is believed preferable to present it as part of a pharmaceutical formulation. In accordance with this aspect of the invention, the pharmaceutical compositions comprise IL-11 in a therapeutically effective dose together with one or more pharmaceutically acceptable carriers and optionally other therapeutic ingredients. A wide variety of pharmaceutically acceptable carriers are
20 known to those of skill in the art. See, e.g., *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 17th ed. (1985), which is incorporated by reference herein. Preferred carriers include inert, non-toxic solids (e.g., dextrose, dextrin, cellulose, pectin, starch, lactose, sucrose, and calcium phosphate,), semi-solids (e.g., glycerol stearate, polyethylene glycol, stearic acid, agar, gelatin, and propylene glycol)
25 and liquids (e.g., buffered saline, water, an organic solvent, and pharmaceutically acceptable oils or fats).

As will be appreciated by those of skill in the art, the preferred form of the pharmaceutical composition of IL-11 will depend on the intended mode of

administration, which in turn will depend on the location and nature of the inflammatory disorder to be treated. For example, delivery to the mouth (e.g., for treatment of oral mucositis, aphthous ulcers, gingivitis, periodontitis), head and/or neck (e.g., for treatment of pharyngitis, esophagitis) can be in the form of aqueous-based oral solutions, suspensions, emulsions, syrups, elixirs, gels, patches, lozenges, tablets, or capsules. Delivery to the gastrointestinal tract (e.g., for treatment of gastrointestinal mucositis, peptic ulcers and inflammatory bowel diseases, such as Crohn's disease, ulcerative colitis, indeterminate colitis, and infectious colitis) can be in the form of oral solutions, gels, suspensions, tablets, capsules, and the like. IL-11, either alone or in combination with other components in a pharmaceutical composition, can be added to infant formula for delivery to the gastrointestinal tract of an infant suffering from or susceptible to necrotizing enterocolitis. It is also possible to formulate the IL-11 preparation for rectal administration (e.g., for treatment of proctitis), e.g., in the form of enema, suppositories, rectal-foam, and the like. Delivery to the eye (e.g., for treatment of ocular diseases such as conjunctivitis, retinitis, and uveitis) can be in the form of solutions, gels, or suspensions. Delivery to the nose (e.g., for treatment of nasal mucositis) can be in the form of solutions, gels, or suspensions. The intranasal formulations may be formulated, for example, into an aqueous or partially aqueous solution, which can then be utilized in the form of a nasal drop or an aerosol. Delivery to the skin (e.g., for treatment of psoriasis) can be in the form of aqueous-based solutions, gels, suspensions, lotions, creams, ointments, patches, and the like.

Liquid carriers are used in preparing solutions, suspensions, emulsions, syrups, elixirs and pressurized compositions. The active ingredient can be dissolved or suspended in a pharmaceutically acceptable liquid carrier such as water, an organic solvent, a mixture of both or pharmaceutically acceptable oils or fats. The liquid carrier can contain other suitable pharmaceutical additives such as solubilizers, emulsifiers, buffers, preservatives, sweeteners, flavoring agents, suspending agents, thickening agents, colors, viscosity regulators, stabilizers or osmo-regulators. Suitable examples of

liquid carriers for oral administration include water (partially containing additives as above), alcohols (including monohydric alcohols and polyhydric alcohols, e.g., glycols) and their derivatives, oils (e.g., peanut oil, sesame oil, olive oil, and coconut oil), and combinations of the above. Compositions comprising such carriers and adjuvants may
5 be formulated using well known conventional materials and methods. Such materials and methods are described, for example, in *Remington's Pharmaceutical Sciences*, *supra*. Infant formula is also a suitable liquid carrier, particularly when the pharmaceutical composition is used to treat or prevent necrotizing enterocolitis.

A solid carrier can include one or more substances which may also act as
10 flavoring agents, lubricants, solubilizers, suspending agents, lubricants, solubilizers, suspending agents, fillers, glidants, compression aids, binders or tablet-disintegrating agents; it can also be an encapsulating material. In powders, the carrier is a finely divided solid which is in admixture with the finely divided active ingredient. In tablets, the active ingredient is mixed with a carrier having the necessary compression
15 properties in suitable proportions and compacted in the shape and size desired. The powders and tablet preferably contain up to 99% of the active ingredient, and may be formulated for immediate and/or sustained release of the active ingredient. Suitable solid carriers include, for example, calcium or sodium phosphate, magnesium stearate, talc, sugars, glycine, lactose, dextrin, starch, gelatin, cellulose, cellulose derivatives
20 (e.g., methyl cellulose, hydroxypropylmethyl cellulose, and sodium carboxymethyl cellulose), polyvinylpyrrolidone, low melting point waxes, and combinations of the above.

Oral tablets may be prepared using a variety of well known methods and in a variety of conventional forms. Exemplary forms include dry powder compaction
25 tablets, micro-particulate systems (e.g., wherein the active ingredient is spray-dried onto a scaffold particle), and hard or soft-gel capsules. The tablets may be optionally covered with an enteric coating, which remains intact in the stomach, but will dissolve and release the contents of the tablet once it reaches the small intestine. Most currently

used enteric coatings are those which remain undissociated in the low pH environment of the stomach, but readily solubilize when the pH rises to about 4 or 5. A number of commercially available enteric coatings may be used depending on the target part of the intestinal tract, i.e., the site of the inflammatory bowel disorder in the patient. Such

5 coatings include, for example, methacrylic acid-methacrylic acid ester-based copolymer, which is sold under the trade name "Eudragit"; anionic water-soluble, polymer cellulose ether, which is sold under the trade name "Aqualon"; cellulose acetate phthalate; polyvinyl acetate phthalate; hydroxypropyl methylcellulose phthalate; and the like. Compositions comprising such carriers and adjuvants may be formulated, and tablets

10 prepared from such compositions, using well known conventional materials and methods. Such materials and methods are described, for example, in *Remington's Pharmaceutical Sciences, supra*.

In one embodiment of the invention, the pharmaceutical composition comprises one or more sustained or controlled release excipients such that a slow or sustained,

15 preferably constant, release of the active ingredient is achieved. A wide variety of suitable excipients are known in the art and are not a part of this invention. Such sustained/controlled release excipients and systems are described, for example, in U.S. Patent No. 5,612,053 (Baichwal et al.), U.S. Patent No. 5,554,387 (Baichwal), U.S. Patent No. 5,512,297 (Baichwal), U.S. Patent No. 5,478,574 (Baichwal et al.), and U.S.

20 Patent No. 5,472,711 (Baichwal et al.), each of which is incorporated by reference herein. If desired, the pharmaceutical composition can be formulated to provide a pulse dose of the active ingredient. A variety of pulse-dose systems, which provide low or high-pulsed doses, are known in the art and are not a part of this invention. Such pulse-dose formulations are described, for example, in Y.H. Bae, "Stimuli-Sensitive Drug

25 Delivery," *Controlled Drug Delivery* (ed. K. Park) ACS Press, Washington, D.C. (1997); *Drugs* (1982) **23**:207-226; *Fertil.Steril.* (1983) **39**:695-699, 1983; *Brit. J. Cancer* (1982) **45**:86-94; *Int. J. Radiat. Oncol. Biol. Phys.* (1982) **8**:915-919; *J. Clin. Endocrinol. Metab.* (1981) **53**:184-91; and *Diabetes* (1977) **26**:571-581.

In another embodiment of the invention, the pharmaceutical composition is formulated to provide direct and/or targeted delivery of the active ingredient to a specific anatomic site or sites within the gastrointestinal tract; e.g., the duodenum, jejunum, ileum, cecum and/or colon, each of which is believed to contain receptors for

- 5 IL-11. Methods for providing targeted delivery of macromolecules, including proteins and polypeptides, to specific tissues or organs within a mammalian host are well known in the art and not a part of this invention. Such targeted delivery systems are described, for example, in R.J. Mersny, "Site-Specific Drug Delivery in the Gastrointestinal Tract," *Controlled Drug Delivery, supra*; M. Mezei and D. Meisner, "Liposomes and
- 10 Nanoparticles as Ocular Drug Delivery Systems," *Biopharmaceutics of Ocular Drug Delivery*, Ch. 6 (ed. P. Edman) CRC Press, Boca Raton, FL (1992); T.L. Bowersock and H. Hogenesch, "Oral Immunization Using Microparticles," *Controlled Drug Delivery, supra*; M. Mezei, "Liposomes in Topical Application of Drugs," *Liposomes as Drug Carriers: Trends and Progress* (ed. G. Gregoriadis), J. Wiley & Sons Publ. USA
- 15 (1988), pp. 663-677; and J.J. Berti and J.J. Lipsky, "Transcutaneous Drug Delivery: A Practical Review," *Mayo CLW Proc.* (1995) 70:581-586.

IL-11, either alone or in combination with other therapeutic agents, may also be administered topically in the form of a dermal patch or transdermal delivery system. Such patches and systems are especially suited for treatment of inflammatory skin

20 disorders such as psoriasis, and for inflammatory disorders involving the mucosa of the head, neck and/or mouth, e.g., oral mucositis, pharyngitis, esophagitis, gingivitis, periodontitis, and aphthous ulcers. In this embodiment of the invention, the pharmaceutical composition may be administered through the use of a dermal patch containing the active ingredient(s) and a carrier that is inert to the active ingredient(s),

25 non-toxic to the skin or mucosal epithelium, and allows delivery of the agent to the dermis and/or epithelium. Dermal patches and delivery systems, utilizing active or passive transdermal delivery carriers, comprising IL-11 may be prepared using well known methods and materials, including, for example, microporous membranes, silicon

polymers and diffusion matrixes. Such materials and methods are described, for example, in *Remington's Pharmaceutical Sciences, supra*.

In yet another embodiment of the invention, at least one proteinase inhibitor is combined with IL-11 in a therapeutic composition to treat inflammatory diseases. In accordance with this aspect of the invention, the pharmaceutical composition comprises IL-11 and one or more proteinase inhibitors, which stabilize the IL-11 molecule against proteolytic degradation. It may be preferable to include a proteinase inhibitor in the pharmaceutical composition to protect IL-11 against proteolytic enzymes in the gastrointestinal tract, particularly in rectal formulations. A wide variety of suitable proteinase inhibitors are known in the art and are not a part of this invention. Suitable proteinase inhibitors include, for example, aprotinin, a kallikrein-trypsin inhibitor, and protein serine proteinase inhibitors such as α -macroglobulin, soybean trypsin inhibitor, and ovomucoid.

A suitable treatment regimen for patients undergoing treatment, including for example prophylactic treatment, may be determined by the attending physician based upon such factors as the patient's age, sex, weight, and general health. Generally, a suitable dose of IL-11 ranges broadly, preferably between about 1 and about 250 microgram (μ g) per kilogram (kg) of body weight of recipient per treatment. Another suitable dose may be in the range of about 1 to about 100 μ g per kg of body weight, and more preferably in the range of about 10 to about 50 μ g per kg of body weight. If desired, these doses can be adjusted to units. A unit is conventionally described as the concentration of polypeptide which leads to half-maximal stimulation in a suitable assay, *e.g.*, for IL-11, the T1165 assay described in PCT/US90/06803. Doses may be administered daily for between one day and six months, or for as long as is deemed necessary and safe, as is readily ascertained by standard tests by the attending physician, depending upon the nature of the disorder being treated. Where appropriate, the dosages may be adjusted upward or downward, for example, a dosing regimen requiring administration of IL-11 at a dose of 25 μ g/kg, daily for one week, or fewer days, or

multiple weeks if indicated. The progress of treatment is appropriately monitored by measurement of markers associated with the disorder being treated to determine if such a dose results in a decrease of for example, TNF- α levels (or corresponding marker) and if not, increasing the dose two-fold for an additional time period of treatment and
5 measurement of marker levels until an effective dosing regimen is reached.

The following examples illustrate the methods of the present invention and in particular the use of topical IL-11 formulations in treating Crohn's disease, chemotherapy-induced ulcerative mucositis, colitis (colonic ulcers), psoriasis, and inflammatory bowel disease. However, as the skilled artisan will recognize from the
10 present disclosure, numerous modifications and variations are possible and comprised within the present invention. The examples are not exhaustive, and do not limit the scope of the invention in any way.

EXAMPLE 1: TREATMENT OF INFLAMMATORY BOWEL DISEASE

In two well-established models it has been shown that subcutaneous rhIL-11 is
15 able to significantly reduce the clinical signs and histologic lesions of gastrointestinal inflammation: acetic acid-induced colitis, a model of acute injury; and the HLA-B27 transgenic rat, a model of chronic inflammation. While conducting further studies of the mechanism of rhIL-11 action, oral topical activity was seen in hamsters with 5-FU induced oral mucositis and mucosal activity was documented in *in vitro/ ex vivo* studies
20 of intestinal transport. Subsequently, these models have been used to confirm the effectiveness of topically and orally administered rhIL-11.

Applicants first examined local topical delivery of rhIL-11 to the colon by enema. To determine the stability of rhIL-11 in the lumen of the gut, rhIL-11 and intestinal chyme from HLA-B27 rats were combined and incubated at 37°C for up to
25 several hours in the presence or absence of a proteinase inhibitors. Fluid samples were analyzed periodically by SDS-PAGE. The addition of a proteinase inhibitor, avian ovinhibitor, stabilized rhIL-11 against the proteolytic degradation observed in the rat

chyme. Next, twenty-three HLA-B27 rats (22-23 weeks old) were randomized to one of six treatment groups: rhIL-11 Vehicle; 50µg/ml ovoinhibitor; rhIL-11 200 µg/ml; or rhIL-11 at 50, 200 or 800 µg/ml combined with 50µg/ml of ovoinhibitor. Animals received a colonic irrigation (1ml) of the respective formulations, Monday, Wednesday and Friday, for three weeks. Stool character was assessed daily. At the end of the experiment, terminal blood samples were obtained, and then the animals were euthanized by an overdose of anesthetic. Platelet counts and colonic mucosal lesions were quantitated and analyzed as described previously. Platelet counts were not different in any of the groups. In the vehicle treated animals, normal, soft or diarrhea stools were present at 16.5, 33, or 50% of the total animal observation days, respectively. In the 50µg/ml+50µg/ml ovoinhibitor animals, normal, soft, or diarrhea stools were present at 65.9, 25, or 9.1% of the total animal observation days, respectively. Histologic lesion scores were 7.25 ± 1.4 and 7.8 ± 1.4 in the vehicle and ovoinhibitor groups, respectively. ANOVA testing showed that all of the rhIL-11 lesion scores were lower than either vehicle or ovoinhibitor ($P < 0.05$), and the lowest lesion score was in the 50µg rhIL-11 + 50µg ovoinhibitor group, 2.1 ± 0.4 . Topically administered rhIL-11 was beneficial in the model of chronic inflammatory bowel disease when combined with a proteinase inhibitor. Since this study was based around a dose of 200µg/ml, no dosing group of 50µg/ml without the proteinase inhibitor was included.

Next, the effects of lower doses of recombinant human interleukin eleven (rhIL-11) with or without Aprotinin (an FDA approved and marketed Proteinase inhibitor) delivered by enema were assessed in the acetic acid model of acute colitis. Colitis was induced in the descending colons of 80 anesthetized Sprague-Dawley rats by intraluminal instillation of 10% acetic acid for 10 seconds on Day 0 of the study. The rats were randomly assigned to the 8 groups according to their different treatments as described below:

TABLE 1

Group	No. Animals	Dosage	Schedule
1 Vehicle	10	1ml.Buffer	3 days/wk/2wks.
2 Aprotinin	10	50µg/ml	3 days/wk/2wks.
5 3 & 4 rhIL-11 ± Aprotinin	20	10µg/µg/ml	3 days/wk/2wks
5 & 6 rhIL-11 ± Aprotinin	20	25µg/g/ml	3 days/wk/2wks
10 7 & 8 rhIL-11 ± Aprotinin	20	50µg/ml	3 days/wk/2wks

Peripheral hematologic parameters were assessed. On day 15, the animals were euthanized, and colons were removed for determination of lesion scores by gross examination and histologic evaluation. Platelet counts in all groups were in normal range. All animals treated with acetic acid developed diffuse colonic lesions, characterized by ulceration, hemorrhage, edema, depletion of goblet cells, and infiltration of leukocytes. The gross and histologic lesions of all rhIL-11 treated animals were significantly reduced compared with vehicle controls on day 15. Administration of rhIL-11 10µg with 50µg Aprotinin 3 days/week/for 2 weeks or 50µg/ml rhIL-11 3 days/wk/2wks appeared to be most effective in this model of colonic mucosal injury.

EXAMPLE 2: TREATMENT OF ORAL MUCOSITIS

95 male Golden Syrian hamsters were equally divided into five groups. Group 1 [control] received phosphate buffered saline and 0.5% hamster serum (vehicle) topically on days 3-14. Groups 2 and 4 received 50 µg IL-11 subcutaneously, twice daily, on days 0-14 and 3-14, respectively. Groups 3 and 5 received 100 µg IL-11 topically, four

times daily, on days 3-14 and 0-14, respectively. Ulcerative mucositis was induced by administration of 60 mg/kg body weight of 5-fluorouracil on days 0 and 2. The left buccal pouch was superficially irritated on day 4. Ulcerative mucositis was evaluated blindly starting on day 6 by scoring standardized photographs. The scoring was done
5 on a scale of 0-10, with 10 being the most severe. Animals were weighed daily. Blood was taken from 3 animals per group on days 6, 10 and 14.

For the entire experiment, average mean ulcerative mucositis scores for all groups were significantly [$p < 0.05$] lower than control. [G1=5.0; G2=3.8; G3=4.1; G4=3.3; G5=4.0]. Day 14 survival rates for G1 through G5 were 23%; 69%, 38%,
10 100% and 46%, respectively. Thus, topical administration of IL-11 was shown to favorably modify the course of chemotherapy-induced ulcerative mucositis relative to control.

The present investigation demonstrates the beneficial effects of a topical formulation of recombinant human IL-11 (rhIL-11) on the induction, course and
15 resolution of chemotherapy-induced mucositis.

Materials and Methods

Male LVG Golden Syrian hamsters (Charles River Laboratories, Wilmington, MA), aged 5-6 weeks, were caged in small groups and fed standard hamster chow and water *ad libitum*. All procedures were conducted in accordance with guidelines set by
20 the Harvard Medical Area Standing Committee on Animals.

Mucositis induction and evaluation

All procedures were performed using anesthesia induced by diethyl ether. To induce mucositis, 5-fluorouracil was administered (60 mg/kg) by intraperitoneal injection on days 0 and 4. To mimic the functional trauma, which occurs in humans, the
25 left buccal pouch mucosa was superficially irritated on day 4. Animals were weighed and observed daily through day 14. Mucositis was evaluated on days 6 to 14. To

evaluate mucositis, animals were anesthetized and the cheek pouch was everted, immobilized, and photographed using a standard, validated technique. At the conclusion of the study, all film was developed simultaneously. Photographs were numbered, randomized and graded blindly by three independent observers using a 10-
 5 point scale in which mild mucositis equaled a score of 1 to 3, moderate 4 to 6, and severe greater than 6. Mild mucositis was defined as mucosal erythema with vasodilation. Mild mucositis was defined as ulceration with pseudomembrane formation, erythema and vasodilation, and severe mucositis consisted of diffuse ulceration with extensive areas of necrosis, pseudomembranes, erythema and
 10 vasodilation.

Test groups and protocol

Ninety-five animals were prospectively randomized into five equal groups. Each animal was individually numbered using an ear punch technique. The following groups were studied:

- | | | |
|----|---------|---|
| 15 | Group 1 | Vehicle control, topical, four times daily, days 3 to 14 (day 0 = day of first chemotherapy injection). |
| | Group 2 | 100 μ g IL-11 per day in a divided dose given twice daily, subcutaneously, days 0-14. |
| 20 | Group 3 | 400 μ g IL-11 per day in a divided dose given four times daily, topically, days 3-14. |
| | Group 4 | 100 μ g IL-11 per day in a divided dose given twice daily, subcutaneously, days 3-14 |
| | Group 5 | 400 μ g IL-11 per day in a divided dose given four times daily, topically, days 0-14. |

25 Animals in the Group 1 were treated with topical application of PBS containing 0.5% hamster serum. Animals in the test groups received rhIL-11 made up in PBS containing 0.5% hamster serum. Animals receiving topical therapy received treatment

at 9 a.m., 11:30 a.m., 2 p.m. and 4:30 p.m. Animals receiving injections of IL-11 were treated at 9 a.m. and 4 p.m. Whole blood was obtained on days 6 and 14 to evaluate the number of platelets.

Results

5 Cumulative mean mucositis scores for group 2 (3.0) and group 4 (2.7) were markedly less than that noted for the control (4.7). Topical application of rhIL-11 was of benefit (group 3 = 3.3; group 5 = 3.4), though slightly less efficacious than rhIL-11 delivered by the subcutaneous route. Interestingly, the efficacy of rhIL-11 given by
10 from day 6 through day 14 among animals treated via the subcutaneous route. On days 6, 7, and 8, animals in group 2 (IL-11 from day 0) had slightly less mucositis than did animals in group 4 (IL-11 beginning on day 3). On day 9 this trend was reversed. No difference was seen for the remaining days of the study.

Schedule of rhIL-11 administration had no observed effect on its topical
15 efficacy, with no significant differences observed between scores of animals in groups 3 and 5. Less mucositis was observed in animals treated with topically applied rhIL-11 throughout the course of the experiment. Reduction in mucositis scores was essentially equivalent between both topical groups. Interestingly, the effect of topically applied rhIL-11 was the same as the subcutaneously administered peptide from day 11 to the
20 conclusion of the experiment.

Evaluation of the effect of rhIL-11 on moderate to severe mucositis demonstrated a similar trend. From day 6 to day 9, the percentage of control animals (group 1) with moderate to severe mucositis was markedly greater than that noted in animals treated with rhIL-11. In general, animals treated subcutaneously responded
25 more favorably. For example, on day 8, 67% of control animals had mucositis scores equal to or greater than 5 compared to 13% of animals in group 2 and 31 % of animals in group 4. The frequency of animals in the topically treated groups was 42% (group 3)

and 39% (group 5). This trend appears to have shifted on days 10 to 11 where it appears that topically treated animals did better than those injected with rhIL-11. While this is true of surviving animals, evaluation of survival data is of importance in interpreting this apparent trend.

- 5 The above results demonstrate that topical administration of rhIL-11 is beneficial *in vivo* for the treatment of oral mucositis.

EXAMPLE 3: TREATMENT OF COLITIS (COLONIC ULCERS)

- 10 The effects of three different dosages of recombinant human interleukin-11 (rhIL-11), given subcutaneously (SC) either prior to or subsequent to intracolonic administration of trinitrobenzene sulfonic acid (TNB), were studied in Sprague-Dawley rats. The TNB or control were given in a 40% ethanol solution to 312 anesthetized adult male rats allotted to one of 26 groups (n=12). Control groups were: subcutaneous (SC); saline alone; intrarectal (IR); 40% ethanol alone; TNB alone; 40% ethanol alone, and SC, rhIL-11 at the highest dosage alone and groups combining TNB with rhIL-11
- 15 therapy, testing three dosages (100, 300, and 1,000 µg/kg), given either before or after induction of colitis with TNB. Body weight changes were monitored. Rats were euthanized at 3 days, 7 days, or 14 days after TNB administration. At necropsy, samples were collected to evaluate fecal occult blood, mucosal myeloperoxidase activity and mucosal gross indexes of ulceration. Histopathologic and ultrastructural analyses of
- 20 the colonic mucosa were performed. The TNB alone elicited a prolonged, severe colitis in treated animals, and the ethanol control group showed a short-lasting, less severe colonic inflammatory response. Colonic ulcer indexes of rhIL-11 treated rats showed a consistent, dose-related reduction in the severity of the TNB-induced colitis, whether the interleukin was given before or after the TNB. This reduction was significant
- 25 (P<0.05) after administration of the intermediate (300 µg/kg) and highest (1,000 µg/kg) dose levels of rhIL-11, in the groups given rhIL-11 for 7 days after TNB. Myeloperoxidase activity was increased during the TNB-induced colitis and was

reduced by rhIL-11 administration ($P < 0.01$). Fecal occult blood loss increased with colitis and paralleled its severity. rhIL-11 enhanced mucus production and decreased the severity of TNB-induced colitis.

EXAMPLE 4: TREATMENT OF MUCOSITIS

- 5 One of the dose-limiting toxicities of radiation therapy for head and neck cancer is the occurrence of oral mucositis. The effects of topical administration of IL-11 on this effect are examined in conjunction with at least one of several regimens employed for the treatment of advanced head and neck cancer. The specifics of the regimens vary, and may include external beam standard fractionated irradiation, hyperfractionated
- 10 irradiation, or accelerated fractionated irradiation. Various chemotherapy agents may also be employed either sequentially or concurrently with radiotherapy. Chemotherapy agents frequently employed in this setting include 5-fluorouracil (5-FU), cisplatin, carboplatin, hydroxyurea, paclitaxel, and topotecan, either alone or in various combinations.
- 15 One regimen is outlined for the purpose of illustration. The regimen involves hyperfractionated external beam irradiation and concurrent chemotherapy with infusional 5-FU and cisplatin. This regimen has been employed for several years for patients with locally advanced squamous cell carcinoma of the head and neck. Patients receive 125 centigray (cG) twice per week of external beam irradiation 5 days per
- 20 week. The total planned dose is 7000 cG. During weeks 1 and 6 of radiotherapy, patients are treated with 5-FU and cisplatin. 5-FU is administered as a 5-day continuous infusion of 600 mg/m²/day while cisplatin is administered as a bolus injection of 12 mg/m²/day for 5 days. There is a 7-day scheduled treatment break after 4000 cG to allow mucositis to subside. The entire planned time for completion of the course of
- 25 irradiation and concurrent chemotherapy is 7 weeks.

Patients are randomly assigned in a ratio of 2:1 to receive either topical IL-11 or placebo. Patients receiving IL-11 will receive either 50 µg/L or 150 µg/mL in 15 to 30

mL of Brioschi (sodium bicarbonate). A 5 mg vial of IL-11 or excipient alone [placebo] first will be resuspended with 1 mL water for injection (WFI). When 30 mL of Brioschi is used, 300 µL (for 50 ug/mL dose) or 950 µL (for the 150 ug/mL dose) containing either IL-11 or excipient alone will be removed from the vial and added to 30 mL

- 5 Brioschi. This final preparation, containing either IL-11 or placebo excipient alone, is swished and swallowed twice daily for 6 to 8 weeks beginning the day before radiotherapy. All patients are assessed to determine the effects of the topical preparation of IL-11 on patients receiving chemotherapy and/or radiotherapy for head and neck cancer. Among the endpoints summarized are:
- 10 • Incidence of oral mucositis of Grade 3 or higher, as measured by the modified National Cancer Institute Common Toxicity Criteria;
 - Duration of oral mucositis of Grade 3 or higher. Duration is measured as (1) the time from onset of Grade 3 or higher until resolution to less than Grade 3 and
 - 15 (2) the number of days of oral mucositis of Grade 3 or higher;
 - Total duration of oral mucositis of any grade;
 - Maximum severity of oral mucositis experienced during the study;
 - Incidence of hospitalization due to oral mucositis; and
 - Incidence of IV fluid resuscitation due to dehydration secondary to poor
 - 20 oral intake [secondary to mucositis].

Patients receiving IL-11 exhibit significant improvement in one or more of the above criteria.

EXAMPLE 5: TREATMENT OF PSORIASIS

- 25 The effects of IL-11 in modulating keratinocyte (the target cell type in a number of cutaneous inflammatory skin diseases, including psoriasis) growth and gene expression were assessed in vitro. When rhIL-11 (recombinant human IL-11) was added to a basal culture medium at a concentration of 100 ng/mL, it stimulated

keratinocyte proliferation. RT-PCR analysis showed that keratinocytes grown in basal as well as enriched medium expressed IL-11 receptor α -chain mRNA (IL-11R α). rhIL-11 was able to stimulate phosphorylation of STAT-3 in keratinocytes when grown in basal medium. To assess whether rhIL-11 could affect the levels of surface intercellular adhesion molecule-1 (ICAM-1) induced by IFN- γ and TNF- α , keratinocytes were exposed to IFN- γ and TNF- α for one hour before adding rhIL-11. Addition of rhIL-11 resulted in a significant decrease in the amount of surface ICAM-1. These results show that IL-11 has direct effects on keratinocytes, thereby downregulating cutaneous immune responses.

10 While the present invention has been described in terms of specific methods and compositions, it is understood that variations and modifications will occur to those skilled in the art upon consideration of the present invention.

 Numerous modifications and variations in the invention as described in the above illustrative examples are expected to occur to those skilled in the art and,
15 consequently, only such limitations as appear in the appended claims should be placed thereon. Accordingly, it is intended in the appended claims to cover all such equivalent variations which come within the scope of the invention as claimed.

 The disclosure of all of the publications cited herein are hereby incorporated by reference.

CLAIMS

What is claimed is:

1. A method of treating an inflammatory disorder, comprising administering a topical formulation of a pharmaceutical composition comprising a pharmaceutically effective amount of IL-11.
2. The method of claim 1 wherein the disorder is mucositis.
3. The method of claim 2 wherein the mucositis is oral mucositis.
4. The method of claim 2 wherein the mucositis is gastrointestinal mucositis.
5. The method of claim 1 wherein the disorder is an inflammatory skin disorder.
6. The method of claim 5 wherein the inflammatory skin disorder is psoriasis.
7. The method of claim 1 wherein the disorder is necrotizing enterocolitis.
8. The method of claim 1 wherein the disorder is an ocular disease.
9. The method of claim 8 wherein the ocular disease is selected from the group consisting of conjunctivitis, retinitis and uveitis.
10. The method of claim 1 wherein the disorder is selected from the group consisting of aphthous ulcers, pharyngitis, esophagitis, peptic ulcers, gingivitis, and periodontitis.

11. The method of claim 1 wherein the pharmaceutically effect amount of IL-11 is between about 1 and about 250 $\mu\text{g/kg}$ body weight.
12. The method of claim 1, wherein the pharmaceutical composition comprises a solution containing IL-11 and a suitable liquid carrier.
- 5 13. The method of claim 12, wherein the suitable liquid carrier is selected from the group consisting of water, organic solvents, oils and fats.
14. The method of claim 12, wherein the suitable liquid carrier is sodium bicarbonate.
15. The method of claim 12, wherein the suitable liquid carrier is an infant formula.
- 10 16. The method of claim 1, wherein the pharmaceutical composition comprises an immediate release carrier for immediate release of IL-11 into the oral cavity.
17. The method of claim 16, wherein the immediate release carrier is selected from the group consisting of sugars, glycine, lactose, dextrin, starch, gelatin, cellulose, methyl cellulose, hydroxypropylmethyl cellulose, and sodium
15 carboxymethyl cellulose.
18. The method of claim 16, wherein the topical formulation is selected from the group consisting of an oral gel, tablet or suspension.
19. The method of claim 1, wherein the pharmaceutical composition comprises an immediate release carrier for delivery of IL-11 to the gastrointestinal tract.

20. The method of claim 19, wherein the immediate release carrier is selected from the group consisting of sugars, glycine, lactose, dextrin, starch, gelatin, cellulose, methyl cellulose, hydroxypropylmethyl cellulose, and sodium carboxymethyl cellulose.
- 5 21. The method of claim 19, wherein the topical formulation is a pill, tablet or capsule.
22. The method of claim 1, wherein the pharmaceutical composition comprises a sustained-release carrier for delivery of IL-11 to the oral cavity.
- 10 23. The method of claim 21, wherein the topical formulation is a patch, lozenge or an uncoated tablet.
24. The method of claim 1, wherein the pharmaceutical composition comprises a sustained-release carrier for delivery of IL-11 to the gastrointestinal tract.
25. The method of claim 24, wherein the topical formulation is a pill, tablet or capsule.
- 15 26. The method of claim 1, wherein the pharmaceutical composition comprises an immediate release carrier for delivery of IL-11 for cervical administration.
27. The method of claim 25, wherein the topical formulation is selected from the group consisting of a topical cream, solution, ointment, and gel.
28. The method of claim 1, wherein the pharmaceutical composition comprises a
20 sustained-release carrier for delivery of IL-11 for cervical administration.

29. The method of claim 26, wherein the topical formulation is a topical cream, solution, ointment or gel.
30. The method of claim 1, wherein the pharmaceutical composition comprises an enema preparation of IL-11 and a suitable liquid carrier for delivery to the colon.
- 5 31. The method of claim 1, wherein the pharmaceutical composition comprises a proteinase inhibitor.
32. The method of claim 31, wherein the proteinase inhibitor is selected from the group consisting of aprotinin, α -macroglobulin, soybean trypsin inhibitor, and ovomucoid.
- 10 33. The method of claim 31, wherein the proteinase inhibitor is aprotinin.
34. The method of claim 1, wherein the topical formulation comprises an enteric coating.
35. The method of claim 34, wherein the enteric coating is selected from the group consisting of a methacrylic acid-methacrylic acid ester-based copolymer, an anionic water-soluble polymer cellulose ether, cellulose acetate phthalate, polyvinyl acetate phthalate, and hydroxypropyl methylcellulose phthalate.
- 15 36. A composition for treating an inflammatory disorder, wherein said composition is a topical formulation comprising a pharmaceutically effective amount of IL-11 and a suitable liquid carrier.
- 20

37. The composition of claim 36, wherein said suitable liquid carrier is sodium bicarbonate.
38. The composition of claim 36, wherein said suitable liquid carrier is an infant formula.
- 5 39. A composition for treating an inflammatory disorder, wherein said composition is a topical formulation comprising a pharmaceutically effective amount of IL-11 and a suitable solid carrier.
40. The composition of claim 39, wherein said suitable solid carrier is an immediate release carrier.
- 10 41. The composition of claim 40, wherein said immediate release carrier is selected from the group consisting of sugars, glycine, lactose, dextrin, starch, gelatin, cellulose, methyl cellulose, hydroxypropylmethyl cellulose, and sodium carboxymethyl cellulose.
- 15 42. The composition of claim 39, wherein said suitable solid carrier is a sustained-release carrier.
43. The composition of claim 39, wherein said suitable solid carrier comprises a proteinase inhibitor.
- 20 44. The composition of claim 39, wherein the proteinase inhibitor is selected from the group consisting of aprotinin, α -macroglobulin, soybean trypsin inhibitor, and ovomucoid.

45. The composition of claim 44, wherein the topical formulation comprises an enteric coating.
46. The composition of claim 45, wherein the enteric coating is selected from the group consisting of a methacrylic acid-methacrylic acid ester-based copolymer,
5 an anionic water-soluble polymer cellulose ether, cellulose acetate phthalate, polyvinyl acetate phthalate, and hydroxypropyl methylcellulose phthalate.

11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500 1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770 1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800 1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1819 1820 1821 1822 1823 1824 1825 1826 1827 1828 1829 1830 1831 1832 1833 1834 1835 1836 1837 1838 1839 1840 1841 1842 1843 1844 1845 1846 1847 1848 1849 1850 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860 1861 1862 1863 1864 1865 1866 1867 1868 1869 1870 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890 1891 1892 1893 1894 1895 1896 1897 1898 1899 1900 1901 1902 1903 1904 1905 1906 1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590 2591 2592 2593 2594 2595 2596 2597 2598 2599 2600 2601 2602 2603 2604 2605 2606 2607 2608 2609 2610 2611 2612 2613 2614 2615 2616 2617 2618 2619 2620 2621 2622 2623 2624 2625 2626 2627 2628 2629 2630 2631 2632 2633 2634 2635 2636 2637 2638 2639 2640 2641 2642 2643 2644 2645 2646 2647 2648 2649

TOPICAL FORMULATION FOR DELIVERY OF INTERLEUKIN-11**ABSTRACT OF THE DISCLOSURE**

Provided by the present invention are topical formulations of Interleukin-11 and methods for treating a variety of disorders, including inflammatory bowel diseases (e.g., Crohn's disease, ulcerative colitis, indeterminate colitis, and infectious colitis), mucositis (e.g., oral mucositis, gastrointestinal mucositis, nasal mucositis, and proctitis), necrotizing enterocolitis, inflammatory skin disorders (e.g., psoriasis, atopic dermatitis, and contact hypersensitivity), aphthous ulcers, pharyngitis, esophagitis, peptic ulcers, gingivitis, periodontitis, and ocular diseases (e.g., conjunctivitis, retinitis, and uveitis).

DECLARATION and POWER OF ATTORNEY

As a below-named inventor, I hereby declare that:

COPY

My residence, post office address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

TOPICAL FORMULATIONS FOR DELIVERY OF INTERLEUKIN-11

the specification of which was filed on October 26, 1998 and assigned Serial No. 09/179,026.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information of which I am aware which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, Section 1.56(a).

I hereby claim foreign priority benefits under Title 35, United States Code, Section 119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s)

<u>Number</u>	<u>Country</u>	<u>Filing Date</u>	<u>Priority Claimed</u>
<u>PCT/US96/08059</u>	<u>PCT</u>	<u>30 May 1996</u>	<u>Yes</u>

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, Section 1.56(a) which occurred

between the filing date of the prior application and the national or PCT international filing date of this application:

COPY


Application Serial Nos.	Filing Date	Status
<u>08/892,407</u>	<u>July 15, 1997</u>	<u>Pending</u>
<u>08/495,724</u>	<u>June 27, 1995</u>	<u>U.S. Patent No. 5,679,339</u>

I hereby appoint the following attorneys to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

Ellen J. Kapinos, Reg. No. 32,245; Steven R. Lazar, Reg. No. 32,618; Scott A. Brown, Reg. No. 32,724; Suzanne A. Sprunger, Ph.D. Reg. No. 41,323; Barbara A. Gyure, Reg. No. 34,614; all of **87 CambridgePark Drive, Cambridge, Massachusetts 02140**; and Thomas J. DesRosier, Reg. No. 30,168; Egon E. Berg, Reg. No. 21,117; Ronald W. Alice, Reg. No. 27,609; of **One Campus Drive, Parsippany, New Jersey 07054**; and Rebecca R. Barrett, Reg. No. 35,152; and Steven R. Eck, Reg. No. 36,126; Arnold S. Milowsky, Reg. No. 35,288; Michael R. Nagy, Reg. No. 33,432; Arthur G. Seifert, Reg. No. 28,040; George Tarnowski, Reg. No. 27,472, all of **P.O. Box 8299, Philadelphia, Pennsylvania, 19101**; Robert F. Boswell, Jr., Reg. No. 35,072, of **P.O. Box 26609, Richmond, Virginia 23261-6609**; and Daniel B. Moran, Reg. No. 41,204 of **401 N. Middletown Road, Pearl River, New York, 10965**.

Address all telephone calls to Barbara A. Gyure at Telephone No. (617) 4986-53. **Address all correspondence to Barbara A. Gyure, Senior Patent Attorney, LEGAL AFFAIRS, GENETICS INSTITUTE, INC., 87 CambridgePark Drive, Cambridge, Massachusetts 02140.**

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full Name of First Inventor Nick W. Warne		
City of Residence Andover	State of Country Massachusetts	Country of Citizenship United States of America
Post Office Address 27 Farrwood Drive	City Andover	State or Country Zip Massachusetts 01810
SIGNATURE: (Please sign and date in permanent blue ink) 		Date Signed: <u>Dec. 7, 1998</u>

COPY

Full Name of First Inventor Camille L. Bedrosian		
City of Residence Belmont Hills	State of Country Massachusetts	Country of Citizenship United States of America
Post Office Address 22 Lantern Road	City Belmont Hills	State or Country Zip Massachusetts 02478-1707
SIGNATURE: (Please sign and date in permanent blue ink) <i>Camille L. Bedrosian</i>		Date Signed: <i>December 9, 1998</i>

Full Name of First Inventor James C. Keith, Jr.		
City of Residence Andover	State of Country Massachusetts	Country of Citizenship United States of America
Post Office Address 28 Vine Street	City Andover	State or Country Zip Massachusetts 01810
SIGNATURE: (Please sign and date in permanent blue ink) <i>James C. Keith, Jr.</i>		Date Signed: <i>12/8/98</i>

Full Name of First Inventor Ullrich S. Schwertschlag		
City of Residence Beverly Farms	State of Country Massachusetts	Country of Citizenship United States of America
Post Office Address 3 Stone Ridge Road	City Beverly Farms	State or Country Zip Massachusetts 01915-2173
SIGNATURE: (Please sign and date in permanent blue ink) <i>Ullrich S. Schwertschlag</i>		Date Signed: <i>12.09.98</i>

Full Name of First Inventor Paul F. Schendel		
City of Residence Wayland	State of Country Massachusetts	Country of Citizenship United States of America
Post Office Address 39 Jeffrey Road	City Wayland	State or Country Zip Massachusetts 01778
SIGNATURE: (Please sign and date in permanent blue ink) <i>Paul F. Schendel</i>		Date Signed: <i>12/10/98</i>